



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

fungus made no further progress, except several spots which showed the brownish hyphæ, but no conidia. The plants upon which no sowings were made remained healthy. If the experiment had been made earlier in the season the development of the fungus would have been more rapid. A form of *C. Apii* is quite common on *Pastinaca*, but is quite distinct from *C. Apii* on cultivated celery. In the *Journal of Mycology* (vol. I, p. 37), the form on *Pastinaca* is included under *C. Apii* Fres. Mr. Ellis thinks, however, that the form on *Pastinaca* might be called *C. Pastinacæ* with propriety, as distinct from *C. Apii* Fres.—B. T. GALLOWAY, *Columbia, Mo.*

An American Papaver.—Last summer Mr. John Spence, a florist of Santa Barbara, collected many plants in the high mountain regions of Santa Barbara County, which he submitted to me for examination. Among other novelties were the beautiful orange-colored blossoms of a *Papaver*. In the absence of root or foliage, it was not easy to tell whether it was *P. Rhœas* of Europe or something new; but, judging from the location in which it was obtained, I suspected it might be something new, and sent it to Dr. Gray. On a visit to the same locality later in the season, Mr. Spence obtained ripe seed, and succeeded in raising perfect plants, specimens of which he has sent to Dr. Gray, who decides that it is a new species. Being the first American representative of the genus, Dr. Gray very appropriately christened it *Papaver Californica*. Mr. Spence says the plant is found in quantity where the ground had been burned over the previous season, and that the large masses of brilliant orange flowers could not fail to attract attention. The same plant was found last year, by other individuals, in two different localities in the Santa Ynez mountains, and under the same conditions, on ground that had been burned over the previous year; but we have no report of its having been collected before last year.—MRS. R. F. BINGHAM.

Vermicularia phlogina Fairman (n. sp.)—Perithecia very delicate, minute, superficial, sparingly clothed with bristles of varying length, subhyaline above, darker below, continuously or sparingly septate, 60-100 μ long and 5-7 μ thick, tips (especially the longer ones) obtuse: sporules oblong, fusoid, hyaline, slightly curved, endochrome at length imperfectly divided in the middle, 15-20 \times 2½-3 μ .—Differs from *V. Coptina* Pk. in its smaller, paler perithecia and different bristles.

On leaves of *Phlox divaricata*. Ridgeway, Orleans County, New York, July, 1886.

This *Vermicularia* occurs on the phlox leaves, (generally after flowering of the host), often mixed with a *Cladosporium* (*C. epiphyllum* Nees?) sometimes pure, and at times suspected leaves have only the *Cladosporium* present.—CHARLES E. FAIRMAN, *Lyndonville, New York.*